



Eidgenössische Technische Hochschule Zürich  
Swiss Federal Institute of Technology Zurich

## Swarming of Brownian Agents

Frank Schweitzer

[fschweitzer@ethz.ch](mailto:fschweitzer@ethz.ch)

*in cooperation with:*

*Werner Ebeling, Benno Tilch, Robert Mach*

# What is it about?



Blue Planet: Seas of Life, Discovery Channel 2001

- ... a swarm ... ... and ...
  - “vortex swarming” of self-propelled animals

## Why do physicists care?

- what is physics? “swarm” of gas molecules in a container
- *passive* vs. *active* motion
  - physics: energy from initial conditions or from *thermal fluctuations* ⇒ Brownian motion
  - biology: energy take-up, storage, conversion (metabolism)
- *undirected* vs. *directed* (correlated) motion
  - physics: driven by convection, currents, external fields
  - biology: “intentional” (preferred direction, avoid collisions, ...)


## Our aims:

- Can we find a *minimal model* of biological swarming?  
... reproduce ‘stylized facts’
- Can we obtain an *analytical description* of the phenomenon?
  - ⇒ derive a macroscopic dynamics from microscopic interactions?
  - ⇒ estimate relevant parameter ranges ? (no blind simulations)
  - ⇒ predict sudden changes in dynamic behavior? (phase transitions)
- Can we *bridge the gap* between ‘physics’ and ‘biology’?
  - ... derive common principles ...
  - ... self-organization, non-equilibrium phase transitions, ...

## Passive Motion

- *undirected motion:*  
driven by thermal noise,  
random impacts  
⇒ Brownian motion
- Fluctuation-Dissipation  
Theorem:

$$S = \gamma \frac{k_B T}{m}$$



## Active Motion

- *active Brownian particles*\* with internal energy depot  $e_i(t)$
- equations of motion:


$$\dot{\mathbf{v}} = -\gamma_0 \mathbf{v} + d_2 e(t) \mathbf{v} + \mathcal{F}^{\text{stoch}}(t)$$

$$\dot{e}(t) = s_0 - c e(t) - d(\mathbf{v}) e(t)$$

- friction function:

$$e_0 = \frac{s_0}{c + d_2 v^2}$$

$$g(v^2) = \gamma_0 - \frac{s_0 d_2}{c + d_2 v^2}$$



\* F.S. *Brownian Agents and Active Particles*, Springer Series in Synergetics, 2003

## Equation of Active Motion:

$$\frac{d\mathbf{v}}{dt} = - \left( \gamma_0 - \frac{s_0 d_2}{c + d_2 v^2} \right) \mathbf{v} + \sqrt{2S} \boldsymbol{\xi}(t)$$

**Supercritical take-up of energy:**  $s_0 d_2 > \gamma_0 c$


$$v_0^2 = \frac{s_0}{\gamma_0} - \frac{c}{d_2} > 0 ; \quad v_0^2 \gg \frac{d}{2} k_B T$$

Start Online-Simulation: Motion of Free Swarm


► mean squared displacement:

$$\Delta R^2(t) = 4D_r t$$


$$\begin{aligned} D_r^{\text{eff}} &= \frac{v_0^4}{2S} \\ &= \frac{1}{2D} \left( \frac{s_0}{\gamma_0} - \frac{c}{d_2} \right)^2 \end{aligned}$$




## Velocity Distribution



$$d_2 = 0.07$$



$$d_2 = 0.2$$



$$d_2 = 0.7$$

## Harmonic Swarms

- parabolic interaction potential  $U(\mathbf{r}_i, \mathbf{R}) = \frac{a}{2} (\mathbf{r} - \mathbf{R})^2$
- equation of motion

$$\begin{aligned}\dot{\mathbf{v}}_i &= -g(v_i^2) \mathbf{v}_i - \nabla U(\mathbf{r}) + \sqrt{2S} \boldsymbol{\xi}_i(t) \\ &= -g(v_i^2) \mathbf{v}_i - \frac{a}{N} \sum_{j=1}^N (\mathbf{r}_i - \mathbf{r}_j) + \sqrt{2S} \boldsymbol{\xi}_i(t)\end{aligned}$$


- supercritical take-up of energy:  $s_0 d_2 > c \gamma_0$   
 $\Rightarrow$  *active motion*  $\Rightarrow$  spatial *dispersion* of the swarm
- mutual interaction  $\Rightarrow$  spatial *concentration*
- **Result:** complex coherent motion

Film

## Superposition of two motions:

- motion of the center of mass ( $\dot{\bar{R}}, \dot{\bar{V}}$ )
- motion *relative to* the center of mass
  - $\bar{V} = 0$  : motion of free particles in parabolic potential
- *dependence on initial conditions:*
  - $v_i^2(0) < v_0^2 \Rightarrow$  rotational mode  
 $V^2(t) \rightarrow 0, v^2(t) \rightarrow v_0^2, r_0 = |v_0| a^{-1/2}$
  - $v_i^2(0) > v_0^2 \Rightarrow$  translational mode  
 $V^2(t) \rightarrow v_0^2, v_i^2(0) > v_0^2$

## Swarming of Daphnia





\* Courtesy of Anke Ordemann, Center for Neurodynamics, University of Missouri

## Interaction via Avoidance Potential

$$\mathcal{F}_i^{\text{ext}} = -\nabla U(\mathbf{r}) = -a\mathbf{r} ; \quad \mathcal{F}_i^{\text{int}} = -\sum_{j \neq i} \nabla V(R_i(\mathbf{r}_{ij}, \mathbf{v}_i, \mathbf{v}_j))$$

Simulation: 20 agents  
distribution of angular momenta



## Two different modes of swarm motion:

- translation/rotation
  - transitions possible dependent on noise, initial conditions

### ➤ *translational mode:*

particles heading into the same direction,  $V^2(t) \neq 0$

Film: Translation of Swarm

Fish

Bird

### ➤ *rotational mode:*

particles moving around a center,  $V^2(t) \rightarrow 0$


Swarm of fish

Simulation

Fish

## Interaction via local alignment

- local orientation of particle:  $\theta_i(t) = \langle \theta(t-1) \rangle_r + \Delta\theta$   
T. Vicsek et al PRL 75 (1995) 1226




H. Levine et al PRE 63 (2001) 017101

- long-range attraction, short-range repulsion

$$U = \sum_{j \neq i} \left\{ C_a \exp(-|x_i - x_j|/l_a) - C_r \exp(-|x_i - x_j|/l_r) \right\}$$

- alignment  $f_i = v_j \exp(-|x_i - x_j|/l_r)$ , or  $f_i = v_i$
- equation of motion:

$$\dot{v}_i = \alpha f_i - \gamma v_i - \nabla U|_{x_i}$$




## Swarming by means of chemical communication

- bacteria, slime molds, amoebae, social insects ...  
use different kind of chemicals for *local* communication
- spatio-temporal communication field

$$\frac{\partial}{\partial t} h_\theta(\mathbf{r}, t) = \sum_{i=1}^N s_i \delta_{\theta, \theta_i} \delta(\mathbf{r} - \mathbf{r}_i) - k_\theta h_\theta(\mathbf{r}, t) + D_\theta \Delta h_\theta(\mathbf{r}, t)$$

- multi-component scalar field reflects:
  - existence of *memory* (past experience)
  - *exchange of information* with *finite* velocity
  - influence of *spatial distances* between agents  
⇒ *weighted* influence (space, time)

## Example: Foraging Route of Ants



Schematic representation of the complete foraging route of *Pheidole milicida*, a harvesting ant of the southwestern U.S. deserts. Each day tens of thousands of workers move out to the dendritic trail system, disperse singly, and forage for food.

Hölldobler, B. and Möglich, M.: The foraging system of *Pheidole militicida* (Hymenoptera: Formicidae), *Insectes Sociaux* 27/3 (1980) 237-264

## Brownian Agents


- two state variables: position  $r_i$   
internal parameter  $\theta_i \in \{-1, +1\}$  (transitions possible)
- dynamic equation for  $r_i$ :

$$\frac{d\mathbf{r}_i}{dt} = \frac{1}{\gamma_0} \nabla_{\mathbf{r}} h^e(\mathbf{r}, t) + \sqrt{2D} \boldsymbol{\xi}_i(t)$$

$$\nabla_{\mathbf{r}} h^e(\mathbf{r}, t) = \frac{\theta_i}{2} \left[ (1 + \theta_i) \nabla_{\mathbf{r}} h_{-1}(\mathbf{r}, t) - (1 - \theta_i) \nabla_{\mathbf{r}} h_{+1}(\mathbf{r}, t) \right]$$

- state dependent production rate  $s_i(\theta_i, t)$

## Non-linear feedback:



➤ **Result:** exploitation of food sources

Film

## Conclusions

- *physical* models of swarm dynamics
  - based on generalized Langevin equations
    - ⇒ derivation of macroscopic dynamics for distributions
    - ⇒ stability analysis: prediction of critical parameters, ....
- turn passive into *active motion*
  - take-off, storage, conversion of energy
- local or global interactions ⇒ *coherent motion*
  - coupling to invariants of motion or local averages
  - attractive, repulsive interaction
  - chemical communication
- *Result*: translational / rotational modes of swarming  
“meaningful behavior”